

ON EIGENVALUES OF THE SCHRÖDINGER OPERATOR WITH AN EVEN COMPLEX-VALUED POLYNOMIAL POTENTIAL

PER ALEXANDERSSON

ABSTRACT. In this paper, we generalize several results of the article “Analytic continuation of eigenvalues of a quartic oscillator” of A. Eremenko and A. Gabrielov.

We consider a family of eigenvalue problems for a Schrödinger equation with even polynomial potentials of arbitrary degree d with complex coefficients, and $k < (d + 2)/2$ boundary conditions. We show that the spectral determinant in this case consists of two components, containing even and odd eigenvalues respectively.

In the case with $k = (d + 2)/2$ boundary conditions, we show that the corresponding parameter space consists of infinitely many connected components.

CONTENTS

1. Introduction	1
1.1. Previous results	2
1.2. Acknowledgements	2
2. Preliminaries on general theory of solutions to the Schroedinger equation	3
2.1. Cell decompositions	3
2.2. From cell decompositions to graphs	4
2.3. The standard order of asymptotic values	5
2.4. Properties of graphs and their face labeling	5
2.5. Braid actions on graphs	7
3. Properties of even actions on centrally symmetric graphs	7
3.1. Additional properties for even potential	7
3.2. Even braid actions	8
4. Proving Main Theorem 1	9
5. Illustrating example	13
References	15

1. INTRODUCTION

We study the problem of analytic continuation of eigenvalues of the Schrödinger operator with an even complex-valued polynomial potential,

Date: January 18, 2013.

2000 Mathematics Subject Classification. Primary 34M40, Secondary 34M03, 30D35.

Key words and phrases. Nevanlinna functions, Schroedinger operator.

that is, analytic continuation of $\lambda = \lambda(\alpha)$ in the differential equation

$$(1) \quad -y'' + P_\alpha(z)y = \lambda y,$$

where $\alpha = (\alpha_2, \alpha_4, \dots, \alpha_{d-2})$ and $P_\alpha(z)$ is the even polynomial

$$P_\alpha(z) = z^d + \alpha_{d-2}z^{d-2} + \dots + \alpha_2z^2.$$

The boundary conditions for (1) are as follows: Set $n = d + 2$ and divide the plane into n disjoint open sectors

$$S_j = \{z \in \mathbf{C} \setminus \{0\} : |\arg z - 2\pi j/n| < \pi/n\}, \quad j = 0, 1, 2, \dots, n-1.$$

The index j should be considered mod n . These are the *Stokes sectors* of the equation (1). A solution y of (1) satisfies $y(z) \rightarrow 0$ or $y(z) \rightarrow \infty$ as $z \rightarrow \infty$ along each ray from the origin in S_j , see [Sib75]. The solution y is called *subdominant* in the first case, and *dominant* in the second case.

The main result of this paper is as follows:

Theorem 1. *Let $\nu = d/2 + 1$ and let $J = \{j_1, j_2, \dots, j_{2m}\}$ with $j_{k+m} = j_k + \nu$ and $|j_p - j_q| > 1$ for $p \neq q$. Let Σ be the set of all $(\alpha, \lambda) \in \mathbf{C}^\nu$ for which the equation $-y'' + (P_\alpha - \lambda)y = 0$ has a solution with the boundary conditions*

$$(2) \quad y \text{ is subdominant in } S_j \text{ for all } j \in J$$

where $P_\alpha(z)$ is an even polynomial of degree d . For $m < \nu/2$, Σ consists of two irreducible connected components. For $m = \nu/2$, which can only happen when $d \equiv 2 \pmod{4}$, Σ consists of infinitely many connected components, distinguished by the number of zeros of the corresponding solution to (1).

1.1. Previous results. The first study of analytic continuation of λ in the complex β -plane for the problem

$$-y'' + (\beta z^4 + z^2)y = \lambda y, \quad y(-\infty) = y(\infty) = 0$$

was done by Bender and Wu [BW69], They discovered the connectivity of the sets of odd and even eigenvalues, rigorous results was later proved in [Sim70].

In [EG09a], the even quartic potential $P_a(z) = z^4 + az^2$ and the boundary value problem

$$-y'' + (z^4 + az^2)y = \lambda_a y, \quad y(\infty) = y(-\infty) = 0$$

was considered.

The problem has discrete real spectrum for real a , with $\lambda_1 < \lambda_2 < \dots \rightarrow +\infty$. There are two families of eigenvalues, those with even index and those with odd. If λ_j and λ_k are two eigenvalues in the same family, then λ_k can be obtained from λ_j by analytic continuation in the complex α -plane. Similar results have been found for other potentials, such as the PT-symmetric cubic, where $P_\alpha(z) = (iz^3 + i\alpha z)$, with $y(z) \rightarrow 0$, as $z \rightarrow \pm\infty$ on the real line. See for example [EG09b].

1.2. Acknowledgements. The author would like to thank Andrei Gabrielov for the introduction to this area of research, and for enlightening suggestions and improvements to the text. Great thanks to Boris Shapiro, my advisor.

2. PRELIMINARIES ON GENERAL THEORY OF SOLUTIONS TO THE SCHROEDINGER EQUATION

We will review some properties for the Schrödinger equation with a general polynomial potential. In particular, these properties hold for an even polynomial potential. These properties may also be found in [EG09a, AG10].

The general Schroedinger equation is given by

$$(3) \quad -y'' + P_\alpha(z)y = \lambda y,$$

where $\alpha = (\alpha_1, \alpha_2, \dots, \alpha_{d-1})$ and $P_\alpha(z)$ is the polynomial

$$P_\alpha(z) = z^d + \alpha_{d-1}z^{d-1} + \dots + \alpha_1 z.$$

We have the associated Stokes sectors

$$S_j = \{z \in \mathbb{C} \setminus \{0\} : |\arg z - 2\pi j/n| < \pi/n\}, \quad j = 0, 1, 2, \dots, n-1,$$

where $n = d+2$, and index considered mod n . The boundary conditions to (3) are of the form

$$(4) \quad y \text{ is subdominant in } S_{j_1}, S_{j_2}, \dots, S_{j_k}$$

with $|j_p - j_q| > 1$ for all $p \neq q$.

Notice that any solution $y \neq 0$ of (3) is an entire function, and the ratio $f = y/y_1$ of any two linearly independent solutions of (3) is a meromorphic function with the following properties, (see [Sib75]).


- (I) For any j , there is a solution y of (3) subdominant in the Stokes sector S_j , where y is unique up to multiplication by a non-zero constant.
- (II) For any Stokes sector S_j , we have $f(z) \rightarrow w \in \bar{\mathbb{C}}$ as $z \rightarrow \infty$ along any ray in S_j . This value w is called *the asymptotic value* of f in S_j .
- (III) For any j , the asymptotic values of f in S_j and S_{j+1} (index still taken modulo n) are distinct. Furthermore, f has at least 3 distinct asymptotic values.
- (IV) The asymptotic value of f in S_j is zero if and only if y is subdominant in S_j . We call such sector *subdominant* for f as well. Note that the boundary conditions given in (4) imply that sectors $S_{j_1}, \dots, S_{j_k}$ are subdominant for f when y is an eigenfunction of (3), (4).
- (V) f does not have critical points, hence $f : \mathbb{C} \rightarrow \bar{\mathbb{C}}$ is unramified outside the asymptotic values.
- (VI) The Schwartzian derivative S_f of f given by

$$S_f = \frac{f'''}{f'} - \frac{3}{2} \left(\frac{f''}{f'} \right)^2$$

equals $-2(P_\alpha - \lambda)$. Therefore one can recover P_α and λ from f .

From now on, f denotes the ratio of two linearly independent solutions of (3), (4).

2.1. Cell decompositions. As above, set $n = \deg P + 2$ where P is our polynomial potential and assume that all non-zero asymptotic values of f are distinct and finite. Let w_j be the asymptotic values of f with an arbitrary ordering satisfying the only restriction that if S_j is subdominant, then $w_j =$

FIGURE 1. Permuting w_j and w_{j+} in Ψ_0 .

0. One can denote by w_j the asymptotic value in the Stokes sector S_j , which will be called the *standard order*, see section 2.3.

Consider the cell decomposition Ψ_0 of $\bar{\mathbb{C}}_w$ shown in Fig. 1a. It consists of closed directed loops γ_j starting and ending at ∞ , where the index is considered mod n , and γ_j is defined only if $w_j \neq 0$. The loops γ_j only intersect at ∞ and have no self-intersection other than ∞ . Each loop γ_j contains a single non-zero asymptotic value w_j of f . For example, for even n , the boundary condition $y \rightarrow 0$ as $z \rightarrow \pm\infty$ for $z \in \mathbb{R}$ implies that $w_0 = w_{n/2} = 0$, so there are no loops γ_0 and $\gamma_{n/2}$. We have a natural cyclic order of the asymptotic values, namely the order in which a small circle around ∞ traversed counterclockwise intersects the associated loops γ_j , see Fig. 1a.

We use the same index for the asymptotic values and the loops, so define

$$j_+ = j + k \text{ where } k \in \{1, 2\} \text{ is the smallest integer such that } w_{j+k} \neq 0.$$

Thus, γ_{j_+} is the loop around the next to w_j (in the cyclic order mod n) non-zero asymptotic value. Similarly, γ_{j_-} is the loop around the previous non-zero asymptotic value.

2.2. From cell decompositions to graphs. Proofs of all statements in this subsection can be found in [EG09a].

Given f and Ψ_0 as above, consider the preimage $\Phi_0 = f^{-1}(\Psi_0)$. Then Φ_0 gives a cell decomposition of the plane $\mathbb{C}_z$. Its vertices are the poles of f and the edges are preimages of the loops γ_j . An edge that is a preimage of γ_j is labeled by j and called a j -edge. The edges are directed, their orientation is induced from the orientation of the loops γ_j . Removing all loops of Φ_0 , we obtain an infinite, directed planar graph Γ , without loops. Vertices of Γ are poles of f , each bounded connected component of $\mathbb{C} \setminus \Gamma$ contains one simple zero of f , and each zero of f belongs to one such bounded connected component. There are at most two edges of Γ connecting any two of its vertices. Replacing each such pair of edges with a single undirected edge and making all other edges undirected, we obtain an undirected graph T_Γ .

It has no loops or multiple edges, and the transformation from Φ_0 to T_Γ can be uniquely reversed.

A *junction* is a vertex of Γ (and of T_Γ) at which the degree of T_Γ is at least 3. From now on, Γ refers to both the directed graph without loops and the associated cell decomposition Φ_0 .

2.3. The standard order of asymptotic values. For a potential P of degree d , the graph Γ has $n = d + 2$ infinite branches and n unbounded faces corresponding to the Stokes sectors of P . We fixed earlier the ordering $w_0, w_1, \dots, w_{n-1}$ of the asymptotic values of f .

If each w_j is the asymptotic value in the sector S_j , we say that the asymptotic values have *the standard order* and the corresponding cell decomposition Γ is a *standard graph*.

Lemma 2 (See Prop. 6 [EG09a]). *If a cell decomposition Γ is a standard graph, then the corresponding undirected graph T_Γ is a tree.*

In the next section, we define some actions on Ψ_0 that permute non-zero asymptotic values. Each unbounded face of Γ (and T_Γ) will be labeled by the asymptotic value in the corresponding Stokes sector. For example, labeling an unbounded face corresponding to S_k with w_j or just with the index j , indicates that w_j is the asymptotic value in S_k .

From the definition of the loops γ_j , a face corresponding to a dominant sector has the same label as any edge bounding that face. The label in a face corresponding to a subdominant sector S_k is always k , since the actions defined below only permute non-zero asymptotic values.

An unbounded face of Γ is called (sub)dominant if the corresponding Stokes sector is (sub)dominant.

2.4. Properties of graphs and their face labeling.

Lemma 3 (See Section 3 in [EG09a]). *Any graph Γ have the following properties:*

- (I) *Two bounded faces of Γ cannot have a common edge, (since a j -edge is always at the boundary of an unbounded face labeled j .)*
- (II) *The edges of a bounded face of a graph Γ are directed clockwise, and their labels increase in that order. Therefore, a bounded face of T_Γ can only appear if the order of w_j is non-standard.*
- (III) *Each label appears at most once in the boundary of any bounded face of Γ .*
- (IV) *The unbounded faces of Γ adjacent to a junction u , always have the labels cyclically increasing counterclockwise around u .*
- (V) *The boundary of a dominant face labeled j consists of infinitely many directed j -edges, oriented counterclockwise around the face.*
- (VI) *If $w_j = 0$ there are no j -edges.*
- (VII) *Each vertex of Γ has even degree, since each vertex in $\Phi_0 = f^{-1}(\Psi_0)$ has even degree, and removing loops to obtain Γ preserves this property.*

Following the direction of the j -edges, the first vertex that is connected to an edge labeled j_+ is the vertex where the j -edges and the j_+ -edges *meet*. The last such vertex is where they *separate*. These vertices, if they exist, must be junctions.

Definition 4. Let Γ be a standard graph, and let $j \in \Gamma$ be a junction where the j -edges and j_+ -edges separate. Such junction is called a j -junction.

There can be at most one j -junction in Γ , the existence of two or more such junctions would violate property (III) of the face labeling. However, the same junction can be a j -junction for different values of j .

There are three different types of j -junctions, see Fig. 2.




FIGURE 2. Different types of j -junctions.

Case (a) only appears when $w_{j+1} \neq 0$. Cases (b) and (c) can only appear when $w_{j+1} = 0$. In (c), the j -edges and j_+ -edges meet and separate at different junctions, while in (b), this happens at the same junction.

Definition 5. Let Γ be a standard graph with a j -junction u . A structure at the j -junction is the subgraph Ξ of Γ consisting of the following elements:

- The edges labeled j that appear before u following the j -edges.
- The edges labeled j_+ that appear after u following the j_+ -edges.
- All vertices the above edges are connected to.

If u is as in Fig. 2a, Ξ is called an I -structure at the j -junction. If u is as in Fig. 2b, Ξ is called a V -structure at the j -junction. If u is as in Fig. 2c, Ξ is called a Y -structure at the j -junction.

Since there can be at most one j -junction, there can be at most one structure at the j -junction.

A graph Γ shown in Fig. 3 has one (dotted) I -structure at the 1-junction v , one (dotted) I -structure at the 4-junction u , one (dashed) V -structure at the 2-junction v and one (dotdashed) Y -structure at the 5-junction u .

Note that the Y -structure is the only kind of structure that contains an additional junction. We refer to such additional junctions as Y -junctions. For example, the junction marked y in Fig. 3 is a Y -junction.




FIGURE 3. Graph Γ with (dotted) I -structures, a (dashed) Y -structure and a (dotdashed) Y -structure.

2.5. Braid actions on graphs. As in [AG10], we define continuous deformations A_j of the loops in Fig. 1a, such that the new loops are given in terms of the old ones by

$$A_j(\gamma_k) = \begin{cases} \gamma_j \gamma_{j+} \gamma_j^{-1} & \text{if } k = j \\ \gamma_j & \text{if } k = j_+ \\ \gamma_k & \text{otherwise} \end{cases}, \quad A_j^{-1}(\gamma_k) = \begin{cases} \gamma_{j+} & \text{if } k = j \\ \gamma_{j+}^{-1} \gamma_j \gamma_{j+} & \text{if } k = j_+ \\ \gamma_k & \text{otherwise} \end{cases}$$

These actions, together with their inverses, generate the Hurwitz (or sphere) braid group $\mathcal{H}_m$, where m is the number of non-zero asymptotic values. (For a definition of this group, see [LZ04].) The action of the generators A_j and A_k commute if $|j - k| \geq 2$.


The property (V) of the eigenfunctions implies that each A_j induces a monodromy transformation of the cell decomposition Φ_0 , and of the associated directed graph Γ .

3. PROPERTIES OF EVEN ACTIONS ON CENTRALLY SYMMETRIC GRAPHS

3.1. Additional properties for even potential. In addition to the previous properties for general polynomials, these additional properties holds for even polynomial potentials P (see [EG09a]). From now until the end of the article, $\nu = (\deg(P) + 2)/2$.

Each solution y of (1) is either even or odd and we may choose y and y_1 such that $f = y/y_1$ is odd.

If the asymptotic values $w_0, w_1, \dots, w_{2\nu-1}$ are ordered in the standard order, we have that $w_j = -w_{j+\nu}$.

FIGURE 4. $E_j(\Psi_0)$

We may choose the loops centrally symmetric in Fig. 1a which implies that Φ_0 and Γ are centrally symmetric.

3.2. Even braid actions. Define the *even actions* E_j as $E_j = A_j \circ A_{j+\nu}$.

Assume that Γ is a graph with the property that if w_j is the asymptotic value in S_k , then w_{j+n} is the asymptotic value in $S_{k+\nu}$. (For example, all standard graphs have this property, with $j = k$.) It follows from the symmetric property of E_j that E_j preserves this property. To illustrate, we have that $E_j(\Psi_0)$ is given in Fig. 4.

Lemma 6. *If Γ is centrally symmetric, then $E_j(\Gamma)$ and $E_j^{-1}(\Gamma)$ are centrally symmetric graphs.*

Proof. We may choose the deformations of the paths γ_j and $\gamma_{j+\nu}$ being centrally symmetric, which implies that the composition $A_j \circ A_{j+\nu}$ preserves the property of Γ being centrally symmetric, see details in [EG09a]. $\square$

Lemma 7. *Let Γ be a centrally symmetric standard graph with no j -junction. Then $E_j^2(\Gamma) = \Gamma$.*

Proof. Since A_j and A_{j+n} commute, we have that $E_j^2 = A_j^2 A_{j+\nu}^2$, and the statement then follows from [AG10, Lemma 12]. $\square$

Theorem 8. *Let Γ be a centrally symmetric standard graph with a j -junction u . Then $E_j^2(\Gamma) \neq \Gamma$, and the structure at the j -junction is moved one step in the direction of the j -edges under E_j^2 . The inverse of E_j^2 moves the structure at the j -junction one step backwards along the j_+ -edges.*

Since Γ is centrally symmetric, it also has a $j + \nu$ -junction, and the structure at the $j + \nu$ -junction is moved one step in the direction of the $j + n$ -edges under E_j^2 . The inverse of E_j^2 moves the structure at the $j + \nu$ -junction one step backwards along the $(j + \nu)_+$ -edges.

Proof. Since $E_j^2 = A_j^2 A_{j+\nu}^2$, the result follows from [AG10, Theorem 13]. $\square$

4. PROVING MAIN THEOREM 1

Notice that each centrally symmetric standard graph Γ has either a vertex in its center, or a double edge, connecting two vertices. This property follows from the fact that Γ_T is a centrally symmetric tree.

Lemma 9. *Let Γ be a centrally symmetric graph. Then for every action E_j , Γ has a vertex at the center iff $E_j(\Gamma)$ has a vertex at the center.*

Proof. This is evident from the definition of the actions, since the action only changes the edges, and preserves the vertices. $\square$

Corollary 10. *The spectral determinant has at least two connected components.*

Each centrally symmetric standard graph Γ is of one of two types:

- (1) Γ has a central double edge. The vertices of the central double edge are called *root junctions*.
- (2) Γ has a junction at its center. This junction is called the *root junction* u_r .

Definition 11. *A centrally symmetric standard graph Γ is in ivy form if Γ consists of structures connected to one or two root junctions.*

Definition 12. *Let Γ be a centrally symmetric standard graph.*

The root metric of Γ , denoted $|\Gamma|_r$ is defined as

$$|\Gamma|_r = \sum_{v \in \Gamma} (\deg(v) - 2) |v - u_r|$$

where the sum is taken over all vertices v of Γ_1 . Here $\deg(v)$ is the total degree of the vertex v in T_Γ and $|v - u_r|$ is the length of the shortest path from v to the closest root junction u_r in T_Γ .

Lemma 13. *The graph Γ is in ivy form if and only if all but its root junctions are Y -junctions.*

Proof. This follows from the definitions of the structures. $\square$

Theorem 14. *Let Γ be a centrally symmetric standard graph. Then there is a sequence of even actions $E^* = E_{j_1}^{\pm 2}, E_{j_2}^{\pm 2}, \dots$, such that $E^*(\Gamma)$ is in ivy form.*

Proof. Assume that Γ is not in ivy form.

Let U be the set of junctions in Γ that are not Y -junctions. Since Γ is not in ivy form we have that $|U| \geq 3$. Let $u_r \neq u_1$ be two junctions in U such that $|u_r - u_1|$ is maximal, and u_r is the central junction closest to u_1 . Let p be the path from u_r to u_1 in T_Γ . It is unique since T_Γ is a tree. Let v be the vertex preceding u_1 on the path p . The edge from v to u_1 in T_Γ is adjacent to at least one dominant face with label j such that $w_j \neq 0$. Therefore, there exists a j -edge between v and u_1 in Γ . Suppose first that this j -edge is directed from u_1 to v . Let us show that in this case u_1 must be a j -junction, i.e., the dominant face labeled j_+ is adjacent to u_1 .

Since u_1 is not a Y -junction, there is a dominant face adjacent to u_1 with a label $k \neq j, j_+$. Hence no vertices of p , except possibly u_1 can be adjacent to j_+ -edges. If u_1 is not a j -junction, there are no j_+ -edges adjacent to u_1 . This implies that any vertex of Γ adjacent to a j_+ -edge is further away from u_r than u_1 .

Let u_2 be the closest to u_1 vertex of Γ adjacent to a j_+ -edge. Then u_2 should be a junction of T_Γ , since there are two j_+ -edges adjacent to u_2 in Γ and at least one more vertex (on the path from u_1 to u_2) which is connected to u_2 by edges with labels other than j_+ . Since u_2 is further away from u_r than u_1 and the path p is maximal, u_2 must be a Y -junction. If the j -edges and j_+ -edges would meet at u_2 , u_1 would be a j -junction. Otherwise, a subdominant face labeled $j + 1$ would be adjacent to both u_1 and u_2 , while a subdominant face adjacent to a Y -junction cannot be adjacent to any other junctions.

Hence u_1 must be a j -junction. By Theorem 8, the action E_j^2 moves the structure at the j -junction u_1 one step closer to u_r along the path p , and similarly happens on the opposite side of Γ , decreasing $|\Gamma|_c$ by at least 2.

The case when the j -edge is directed from v to u_1 is treated similarly. In that case, u_1 must be a j_- -junction, and the action $A_{j_-}^{-2}$ moves the structure at the j_- -junction u_1 one step closer to u_r along the path p .

We have proved that if $|U| > 1$ then $|\Gamma|_r$ can be reduced. Since it is a non-negative integer, after finitely many steps we must reach a stage where U consists only of the root junctions. Hence $E^*(\Gamma)$ is in ivy form. $\square$

The above Theorem shows that for every centrally symmetric standard graph Γ , there is a sequence of actions that turns Γ into ivy form. A graph in ivy form consists of one or two root junctions, with attached structures. These structures can be ordered counterclockwise around each root junction. These observations motivates the following lemmas:

Lemma 15. *Let Γ be a centrally symmetric standard graph, and let $u_r \in \Gamma$ be a root junction of type j_- and of type j . Let S_1 and S_2 be the corresponding structures attached to u_r .*

- (1) *If S_1 and S_2 are of type Y resp. V , then there is a sequence of even actions that interchange these structures.*
- (2) *If S_1 and S_2 are of type I resp. Y , then there is a sequence of even actions that converts the type Y structure to a type V structure.*
- (3) *If S_1 and S_2 are both of type Y , then there is a sequence of even actions that converts one of the Y -structures to a V -structure.*

Proof. By symmetry, there are identical structures in Γ attached to a root junction of type $n + j_-$ and $n + j$, with attached structures S'_1 and S'_2 of the same type as S_1 resp. S_2 .

Lemma 19, 20 and 22 in [AG10], gives the existence of a non-even sequence of actions, that only acts on S_1 and S_2 in the desired way.

In all these cases, the sequence is of the form

$$A^* = A_{k_1}^{\pm 2} A_{k_2}^{\pm 2} \dots A_{k_m}^{\pm 2}$$

where $k_1, k_2, \dots, k_m \in \{j, j_-\}$. It follows that the action

$$B^* = A_{k_1+\nu}^{\pm 2} A_{k_2+\nu}^{\pm 2} \dots A_{k_m+\nu}^{\pm 2}$$

do the same as A^* but on S'_1 and S'_2 .

Now, $E^* = A^* \circ B^*$ is even, since by commutativity¹, it is equal to

$$(A_{k_1}^{\pm 2} A_{k_1+\nu}^{\pm 2})(A_{k_2}^{\pm 2} A_{k_2+\nu}^{\pm 2}) \dots (A_{k_m}^{\pm 2} A_{k_m+\nu}^{\pm 2})$$

which easily may be written in terms of our even actions as

$$E_{k_1}^{\pm 2} E_{k_2}^{\pm 2} \dots E_{k_m}^{\pm 2}.$$

This sequence of actions has the desired property. $\square$

Corollary 16. *Let Γ be a centrally symmetric graph, with two adjacent dominant faces. Then there is a sequence of even actions E^* such that $E^*(\Gamma)$ has either one or two junctions.*

Proof. We may apply even actions to make Γ into a standard graph, and then convert it to ivy form. The condition that we have two dominant faces, is equivalent to existence of I -structures. If there are no Y -structures, then the only junctions of Γ are the root junctions, and we are done. Otherwise, we may have the Y - and V -structures, so that a Y -structure appears next to the I -structure. By using the second part of the above lemma, we decrease the number of Y -structures of Γ by two. After a finite number of actions, we arrive at a graph in ivy form without Y -structures. $\square$

Lemma 17. *Let Γ be a centrally symmetric graph, with no adjacent dominant faces. Then there is a sequence of even actions E^* such that $E^*(\Gamma)$ is in ivy form, with at most two Y -structures.*

Proof. By Theorem 14, we may assume that Γ is in ivy form. Since there are no adjacent dominant sectors, the only structures of Γ are of Y and V type. These are attached to the one or two root junctions.

Assume that there are more than two Y -structures present. Two of these must be attached to the same root junction, u_r . By repeatedly applying part one of Lemma 15, we may interchange the Y - and V -structures attached to u_r such that the two Y -structures are adjacent. Applying part three of Lemma 15, we may then convert one of the two Y -structures to a V -structure.

By symmetry, the same change is done on the opposite side of Γ and total number of Y -structures of Γ have therefore been reduced by two. We may repeat this procedure a finite number of times, until the number of Y -structures is less than three. This implies the lemma. $\square$

Lemma 18 (See [AG10]). *Let Γ be a standard graph such that no two dominant faces are adjacent. Then the number of bounded faces of Γ is finite and does not change after any action A_j^2 .*

Corollary 19. *The number of bounded faces of Γ does not change under any even action $E_j^2 = A_j^2 A_{j+\nu}^2$.*

Lemma 20. *Let $\nu = n/2 = d/2 + 1$ and let Σ be the space of all $(\alpha, \lambda) \in \mathbb{C}^{\nu-1}$ such that equation (1) admits a solution subdominant in non-adjacent Stokes sectors*

$$(5) \quad S_{j_1}, S_{j_2}, \dots, S_{2m}$$

¹We have at least 4 structures, 2 of them are Y or V structures. Hence $n \geq 6$ and we have commutativity.

with $j_{k+m} = j_k + \nu$ and $1 \leq m \leq \nu/2$. Then Σ is a smooth complex analytic submanifold of $\mathbf{C}^{\nu-1}$ of the codimension m .

Proof. We consider the space $\mathbf{C}^{\nu-1}$ as a subspace of the space $\mathbf{C}^{n-2}$ of all (α, λ) corresponding to the general polynomial potentials in (3), with $\alpha = (\alpha_1, \dots, \alpha_{d-1})$. Let f be a ratio of two linearly independent solutions of (3), and let $w = (w_0, \dots, w_{n-1})$ be the set of the asymptotic values of f in the Stokes sectors $S_0, \dots, S_{n-1}$.

Then w belongs to the subset Z of $\bar{\mathbf{C}}^{n-1}$ where the values w_j in adjacent Stokes sectors are distinct and there are at least three distinct values among w_j . The group G of fractional-linear transformations of $\bar{\mathbf{C}}$ acts on Z diagonally, and the quotient Z/G is a $(n-3)$ -dimensional complex manifold.

Theorem 7.2, [Bak77] implies that the mapping $W : \mathbf{C}^{n-2} \rightarrow Z/G$ assigning to (α, λ) the equivalence class of w is submersive. More precisely, W is locally invertible on the subset $\{\alpha_{d-1} = 0\}$ of $\mathbf{C}^{n-2}$.

For an even potential, there exists an odd function f . The corresponding set of asymptotic values satisfies ν linear conditions $w_{j+\nu} = -w_j$ for $j = 0, \dots, \nu-1$. For $(\alpha, \lambda) \in \Sigma$, we can assume that $S_{j_1}, \dots, S_{j_m}$ are subdominant sectors for f . This adds m linearly independent conditions $w_{j_1} = \dots = w_{j_m} = 0$. Let Z_0 be the corresponding subset of Z . Its codimension in Z is $\nu + m$. The one-dimensional subgroup $\mathbf{C}^*$ of G consisting of multiplications by non-zero complex numbers preserves Z_0 , and $gZ_0 \cap Z_0 = \emptyset$ for each $g \in G \setminus \mathbf{C}^*$. The explanation is as follows:

Since we have at least two subdominant sectors, only fractional linear transforms that preserves 0 are allowed. Furthermore, there exists a sector S_k with the value w_k different from 0 and ∞ (otherwise we would have only two asymptotic values). There is a unique transformation, multiplication by w_k^{-1} , preserving 0 and sending $\pm w_k$ to ± 1 . This implies that the only transformation preserving 0 and sending $\pm w_k$ to another pair of opposite numbers is multiplication by a non-zero constant.

Hence GZ_0 is a G -invariant submanifold of Z of codimension $\nu + m - 2$, and its image $Y_0 \subset Y$ is a smooth submanifold of codimension $\nu + m - 2$. Due to Bakken's theorem, $W^{-1}(Y_0)$ intersected with the $(n-3)$ -dimensional space of (α, λ) with $\alpha_{d-1} = 0$ is a smooth submanifold of codimension $\nu + m - 2$, dimension $\nu - m - 1$. Accordingly, it is a smooth submanifold of codimension m of the space $\mathbf{C}^{\nu-1}$. $\square$

Proposition 21. *Let Σ be as in Lemma 20. If at least two adjacent Stokes sectors are missing in (5), then Σ consists of two irreducible complex analytic manifolds.*

Proof. Nevanlinna theory (see [Nev32, Nev53]), implies that, for each symmetric standard graph Γ with the properties listed in Lemma 3, there exists $(\alpha, \lambda) \in \mathbf{C}^{n-1}$ and an odd meromorphic function $f(z)$ such that f is the ratio of two linearly independent solutions of (1) with the asymptotic values w_j in the Stokes sectors S_j , and Γ is the graph corresponding to the cell decomposition $\Phi_0 = f^{-1}(\Psi_0)$. This function, and the corresponding point (α, λ) is defined uniquely.

Let $W : \Sigma \rightarrow Y_0$ be as in the proof of Lemma 20. Then Σ is an unramified covering of Y_0 . Its fiber over the equivalence class of $w \in Y_0$ consists of the points $(\alpha_\Gamma, \lambda_\Gamma)$ for all standard graphs Γ . Each action A_j^2 corresponds

to a closed loop in Y_0 starting and ending at w . It should be noted that Y_0 is a connected manifold. Since for a given list of subdominant sectors a standard graph with one vertex is unique, Theorem 15 implies that the monodromy group has two orbits; odd and even eigenfunctions cannot be exchanged by any path in Y_0 , while any odd (even) can be transferred into any other odd (even) eigenfunction by a sequence of $E_k^{\pm 2}$.

Hence Σ consists of two irreducible connected components (see, e.g., [Kho04]). $\square$

This immediately implies Theorem 1, for $m < \nu/2$. The following proposition implies the case where $m = \nu/2$.

Proposition 22. *Let Σ be the space of all $(\alpha, \lambda) \in \mathbf{C}^{\nu-1}$, for even ν , such that equation (1) admits a solution subdominant in every other Stokes sector, that is, in $S_0, S_2, \dots, S_{n-2}$.*

Then irreducible components Σ_k , $k = 0, 1, \dots$ of Σ , which are also its connected components, are in one-to-one correspondence with the sets of centrally symmetric standard graphs with k bounded faces. The corresponding solution of (1) has k zeros and can be represented as $Q(z)e^{\phi(z)}$ where Q is a polynomial of degree k and ϕ a polynomial of degree $(d+2)/2$.

Proof. Let us choose w and Ψ_0 as in the proof of Proposition 21. Repeating the arguments in the proof of Proposition 21, we obtain an unramified covering $W : \Sigma \rightarrow Y_0$ such that its fiber over w consists of the points $(\alpha_\Gamma, \lambda_\Gamma)$ for all standard graphs Γ with the properties listed in Lemma 3.

Since we have no adjacent dominant sectors, Lemma 17 implies that any standard graph Γ can be transformed by the monodromy action to a graph Γ_0 in ivy form with at most two Y -structures attached at the root junction(s) of type j and $j + \nu$.

Lemma 18 implies that Γ and Γ_0 have the same number k of bounded faces. If $k = 0$, the graph Γ_0 is unique. If $k > 0$, the graph Γ_0 is completely determined by k . Hence for each $k = 0, 1, \dots$ there is a unique orbit of the monodromy group action on the fiber of W over w consisting of all standard graphs Γ with k bounded faces. This implies that Σ has one irreducible component for each k .

Since Σ is smooth by Lemma 20, its irreducible components are also its connected components.

Finally, let $f_\Gamma = y/y_1$ where y is an odd solution of (1) subdominant in the Stokes sectors $S_0, S_2, \dots, S_{n-2}$. Then the zeros of f and y are the same, each such zero belongs to a bounded domain of Γ , and each bounded domain of Γ contains a single zero. Hence y has exactly k simple zeros. Let Q be a polynomial of degree k with the same zeros as y . Then y/Q is an entire function of finite order without zeros, hence $y/Q = e^\phi$ where ϕ is a polynomial. Since y/Q is subdominant in $(d+2)/2$ sectors, $\deg \phi = (d+2)/2$. $\square$

5. ILLUSTRATING EXAMPLE

We will now give a small example on how to apply the method given in the previous section, Theorem 14 and Lemma 15. Let Γ be as in Fig. 5a. From subsection 2.4, we have that a dominant face with label j have j -edges

as boundaries. Hence the faces 0 and 4 are subdominant. Also, the direction of the edges are directed counterclockwise in each of the dominant faces. Applying E_1^2 , moves the I -structure at the 1-junction one step to the right,




FIGURE 5. The graphs Γ and $E_1^2(\Gamma)$

following the 1-edges. Similarly, the I -structure at the 5-junction moves one step to the left. Therefore, $E_1^2(\Gamma)$ is given in Fig. 5b. The graph $E_1^2(\Gamma)$ is now in ivy form, it consists of a center junction connected to four I -structures and two Y -structures. We proceed by using the algorithm in Lemma 15, and apply E_1^2 two times more. These steps are given in Fig. 6. The next





FIGURE 6. The graphs $E_1^4(\Gamma)$ and $E_1^6(\Gamma)$

step in the lemma is to move the newly created V -structures to the center junction. We therefore apply E_3^2 two times. These final steps are presented in Fig. 7, and we have reached the unique graph with only one junction.

FIGURE 7. The graphs $E_3^2 E_1^6(\Gamma)$ and $E_3^4 E_1^6(\Gamma)$

REFERENCES

- [AG10] P. Alexandersson and A. Gabrielov. On eigenvalues of the schrödinger operator with a polynomial potential with complex coefficients. 2010.
- [Bak77] I. Bakken. A multiparameter eigenvalue problem in the complex plane. *Amer. J. Math.*, 99(5):1015–1044, 1977.
- [BW69] C. Bender and T. Wu. Anharmonic oscillator. *Phys. Rev. (2)*, 184:1231–1260, 1969.
- [EG09a] A. Eremenko and A. Gabrielov. Analytic continuation of eigenvalues of a quartic oscillator. *Comm. Math. Phys.*, 287(2):431–457, 2009.
- [EG09b] A. Eremenko and A. Gabrielov. Irreducibility of some spectral determinants. 2009. arXiv:0904.1714.
- [Kho04] A. G. Khovanskii. On the solvability and unsolvability of equations in explicit form. (russian). *Uspekhi Mat. Nauk*, 59(4):69–146, 2004. translation in Russian Math. Surveys 59 (2004), no. 4, 661–736.
- [LZ04] S. Lando and A. Zvonkin. *Graphs on Surfaces and Their Applications*. Springer-Verlag, 2004.
- [Nev32] R. Nevanlinna. Über Riemannsche Flächen mit endlich vielen Windungspunkten. *Acta Math.*, 58:295–373, 1932.
- [Nev53] R. Nevanlinna. *Eindeutige analytische Funktionen*. Springer, Berlin, 1953.
- [Sib75] Y. Sibuya. *Global theory of a second order differential equation with a polynomial coefficient*. North-Holland Publishing Co., Amsterdam-Oxford; American Elsevier Publishing Co., Inc., New York, 1975.
- [Sim70] B. Simon. Coupling constant analyticity for the anharmonic oscillator. *Ann. Physics*, 58:76–136, 1970.

DEPARTMENT OF MATHEMATICS, STOCKHOLM UNIVERSITY, SE-106 91, STOCKHOLM, SWEDEN

E-mail address: per@math.su.se